

On the Variety of rational Space Curves

Z. Ran

In this paper we study enumeratively the variety $V_{d,n}$ parametrising irreducible nonsingular rational curves of degree d in $\mathbb{P}^n$, $n \geq 3$. We shall give a recursive procedure which computes two sets of numbers associated to $V_{d,n}$:

- The Schubert degrees $N_{d,n}(a_1, \dots, a_k)$, i.e. the degree of the locus of members of $V_{d,n}$ meeting a generic collection $A_1, \dots, A_k$ of linear subspaces of respective codimensions $a_1, \dots, a_k$ in $\mathbb{P}^n$, whenever $\sum(a_i - 1) = (n+1)d - (n-3) = \dim V_{d,n}$, i.e. whenever the locus in question is finite;
- the linear genera $g_{d,n}(a_1, \dots, a_k)$, i.e. the (geometric) genus of the analogously-defined locus whenever it is 1-dimensional, i.e. whenever $\sum(a_i - 1) = \dim V_{d,n} - 1$.

Analogous questions for $n = 2$ were considered in [1] [2]. As there, the method is completely elementary, involving some geometry on ruled surfaces. No quantum-cohomological methods are used, indeed our method may be viewed as an alternative to those (compare [3] and references therein).

1. Degrees.

In what follows we fix $n \geq 3$ and denote by $\bar{V}_d$ the closure in the Hilbert scheme of the locus of irreducible nonsingular rational curves of degree d in $\mathbb{P}^n$. Let $A_1, \dots, A_k$ be a generic collection of linear subspaces of respective codimensions $a_1, \dots, a_k \geq 1$ in $\mathbb{P}^n$. We denote by

$$B_d = B_d(a.) = B_d(A.)$$

the normalization of the locus

$$\{(C, P_1, \dots, P_k) : C \in \bar{V}_d, P_i \in C \cap A_i, i = 1, \dots, k\}$$

when all $a_i > 1$, this is also the normalization of its projection to $\bar{V}_d$, i.e. the locus of degree- d rational curves (and their specializations) meeting $A_1, \dots, A_k$; however it will be convenient to allow some A_i to be hyperplanes. Of course if some $a_i > n$ then $N_d(a.) = 0$. Let's call the number of i such that $a_i > 1$ the *length* of the condition vector $(a.)$. We have

$$\dim B = (n+1)d - (n-3) - \sum(a_i - 1).$$

When this is 0, we set

$$N_d(a.) = \deg B_d(a.).$$

Of course $N_d(1, a_2, \dots) = dN_d(a_2, \dots)$, so it will suffice to compute these when all $a_i > 1$.

The plan is to get at them via suitable 1-dimensional B 's. To this end, take a $B = B(A.)$ 1-dimensional and let

be the normalization of the tautological family of rational curves, and $f : X \rightarrow \mathbb{P}^n$ the natural map. Then X is smooth and each fibre of π is either a $\mathbb{P}^1$ or a pair of $\mathbb{P}^1$'s meeting transversely once. Let $\mathcal{F}$ be the set of components of reducible fibres of π . Now X comes equipped with a set of distinguished sections $s_i = s_{A_i}, i = 1, \dots, k$ and note that

$$s_i \cdot s_j = N_d(\dots, a_i + a_j, \dots, \hat{a}_j, \dots), i \neq j$$

Also, let $R_i = R_{A_i}$ be the sum of all fibre components not meeting s_{A_i} and $\mathcal{F}_A \subset \mathcal{F}$ the set of such components. Then R_{A_i} may be blown down, giving rise to a geometric ruled surface:

$$(1) \quad b_i : X \rightarrow X_i = X_{A_i} = \mathbb{P}(E_{A_i}),$$

with sections $\bar{s}_j = b_i(s_j)$ and note that if, say, $i = 1$, then

$$\bar{s}_2 \cdot \bar{s}_1 = s_2 \cdot s_1.$$

Now set

$$m_i = m_i(a_1, \dots, a_k) = -s_i^2, i = 1, \dots, k.$$

Note that $m_1 = -\bar{s}_1^2$ too. But clearly, on a geometric ruled surface the difference of any two sections is a sum of fibres hence has self intersection $= 0$, hence

$$\begin{aligned} 0 &= (\bar{s}_1 - \bar{s}_2)^2 \\ &= s_1^2 + s_2^2 + s_2 \cdot R_1 - 2s_2 \cdot s_1, \\ &= -m_1 - m_2 + s_2 \cdot R_1 - 2s_2 \cdot s_1. \end{aligned}$$

i.e.

$$(2) \quad m_1 + m_2 = s_2 \cdot R_1 - 2N_d(a_1 + a_2, a_3, \dots).$$

One consequence of this, already noted and used in [1] is the

2-section lemma 1.1. If $a_1 = a_2$, then

$$(3) \quad s_1^2 = s_2^2 = \frac{-1}{2} s_1 \cdot R_2 + N_d(a_1 + a_2, a_3, \dots)$$

Indeed if $a_1 = a_2$ then clearly by monodromy $m_1 = m_2$ so (3) follows from (2).

For general codimensions we have the

3-section lemma 1.2. For any 3 distinct distinguished sections s_1, s_2, s_3 we have

$$(4) \quad s_1^2 = \frac{-1}{2} (s_1 \cdot R_2 + s_1 \cdot R_3 - s_2 \cdot R_3) + s_1 \cdot s_2 + s_1 \cdot s_3 - s_2 \cdot s_3.$$

This follows immediately from (2) by a suitable linear combination .

Note that by an obvious dimension count the number k of distinguished sections on X is always > 3 , so Lemma 1.2 is always applicable. Also, from a recursive standpoint, numbers such as $s_1 \cdot R_2$, having to do with reducible curves, are easily computable in terms of $N_d, d' < d$, hence may be considered known. Indeed,

$$(5) \quad s_1 \cdot R_2 = \sum N_d(A_1, A_2, \mathbb{P}^{s_1}) N_d(A_2, A_3, \mathbb{P}^{s_2})$$

the summations being over all $d_1 + d_2 = d, s_1 + s_2 = n$ and all decompositions $A. = (A_1, A_2) \coprod (A^1) \coprod (A^2)$ (as unordered sequences or partitions). Each term corresponds to a pair of families of curves each of degree d_i meeting (A_1, A^i) and filling up a locus of codimension s_i and degree $N_{d_i}(A^i, A_i, \mathbb{P}^{s_i})$ in $\mathbb{P}^n$, $i = 1, 2$; the two loci meet in a finite set whose cardinality is given by Bezout's theorem and whose members correspond with $s_1 \cap R_2$.

Thus at least when $a_1 + a_2, a_1 + a_3, a_2 + a_3$ are all $> n$ (which is automatic if $n = 3$ but not otherwise), the 3-section lemma computes m_1, m_2, m_3 in terms of lower-degree data; but even if this condition is not satisfied (and, say, $a_1, a_2, a_3 > 1$), the lemma still computes m_1 , say, in terms of data of lower degree *or lower length*; we shall use this observation below in constructing a 'length recursion'.

Now let $L = f^* \mathcal{O}(1)$ and F_0 be a general fibre of π . Then we have

$$(6) \quad L = ds_1 - \sum_{F \in \mathcal{F}_{A_1}} \deg(F)F + xF_0$$

for some $x \in \mathbb{Q}$: indeed (6) holds simply because both sides have the same value on all fibre components. To determine x , evaluate on s_1 , noting that, by definition,

$$L \cdot s_1 = N_d(a_1 + 1, a_2, \dots).$$

Thus we have

$$x = N_d(a_1 + 1, a_2, \dots) + dm_1.$$

Now let's square (6), noting that, by definition, $L^2 = N_s(2, a_1, \dots)$. Thus

$$N_d(2, a_1, \dots) = -d^2m_1 + 2d^2m_1 + 2dN_d(a_1 + 1, a_2, \dots) - \sum_{F \in \mathcal{F}_{A_1}} (\deg F)^2$$

i.e.

$$(7) \quad N(2, a_1, \dots) = 2dN_d(a_1 + 1, a_2, \dots) + d^2m_1(a.) - \sum_{F \in \mathcal{F}_{A_1}} (\deg F)^2$$

As above, the sum $\sum_{F \in \mathcal{F}_A} (\deg F)^2$ is easily evaluated in terms of $N_{d'}, d' < d$ and may be considered known. In particular, when $a_1 = n$, i.e. A_1 is a point, and moreover $a_2 + a_3 > n$, we have $N_d(a_1 + 1, a_2, \dots) = 0$ and that $m_1(a.)$, via the 3-section lemma, is computable from lower-degree data, so (7) yields a recursive formula for all the $N_d(2, n, a_2, a_3, \dots)$, $a_2 + a_3 > n$, namely

$$(8) \quad N_d(2, n, a_2, \dots) = d^2m_1(n, a_2, \dots) - \sum_{F \in \mathcal{F}_{A_1}} (\deg F)^2$$

Now take the dot product of (6) with s_2 , obtaining

$$(9) \quad N_d(a_1, a_2 + 1, \dots) = dN_d(a_1 + a_2, \dots) - \sum_{F \in \mathcal{F}_{A_1}} (\deg F) + N_d(a_1 + 1, a_2, \dots) + dm_1(a.)$$

Now to determine $N_d(a.)$ in general we proceed by recursion on k (as well as d), as follows. We may assume all $a_i > 1$. For the smallest possible k (given $d \geq 3$), clearly we may assume by reordering that $a_1 = a_2 = a_3 = n$ so from (7) (read backwards) we compute $N_d(a.)$ from $N_d(2, n-1, n, n, \dots) = N_d(2, n, n, n-1, \dots)$ (and lower-degree data); but $N_d(2, n, n, n-1, \dots)$ has already been computed above; this takes care of the case of smallest k . In the general case we use recursion on k . Using (9), we compute $N_d(a.)$ from $N_d(a_1+1, a_2-1, \dots)$ and terms of lower degree or length. Applying (9) repeatedly we compute $N_d(a.)$ in terms of lower-degree and lower-length data plus $N_d(a_1+a_2-1, 1, a_3, \dots) = dN_d(a_1+a_2-1, a_3, \dots)$, itself a lower-length term. This computes $N_d(a.)$ in general.

2. Genera.

In this section we fix a sequence $(a.)$ giving rise to a smooth (maybe disconnected) curve $B = B(a.)$ and give a formula for the latter's geometric genus, i.e. for $\deg(K_B)$. The idea is to consider a 'thickening'

$$B \rightarrow B^+ = B(a^+) = B(A^+), \dim B^+ = n-1,$$

where $A. = (A_1, \dots, A_k)$

$$A^+ = (A_1^+, \dots, A_\ell^+), \ell \leq k$$

$$A. \subseteq A_i^+ \subset \mathbb{P}^n$$

$$a_i^+ = \dim A_i^+ \leq n-2.$$

Consider the diagram (over a neighborhood of the image of B):

$$\begin{array}{ccccc} X & \longrightarrow & X^+ & \xrightarrow{f^+} & \mathbb{P}^n \\ \downarrow & & \downarrow & & \\ B & \longrightarrow & B^+ & & \end{array}$$

with f^+ generically finite. Let ρ be the ramification divisor of f^+ . Then, as in [2] it is easy to see that

$$\rho|_X = \sum (a_i^+ - 1)s_i.$$

On the other hand by Riemann-Hurwitz,

$$(10) \quad -(n+1)L + \rho = K_{X^+}|_X = K_{X/B} + \pi^*(K_{B^+}|_B).$$

Now recall the blowing down map $X \rightarrow X_1 = X_{A_1} = \mathbb{P}(E_{A_1})$ (1). It is easy to see that

$$K_{X_1/B} = -2s_1 - m_1F_0,$$

hence

$$K_{X/B} = -2s_1 - m_1F_0 + R_1$$

so that by (8),

$$\pi^*(K_{B^+}|_B) = -(n+1)L + \sum (a_i^+ - 1)s_i + 2s_1 + m_1F_0 - R_1$$

Evaluating on s_1 , we conclude

$$\deg(K_{B+}|_B) = -(n+1)N_d(a_1+1, a_2, \dots) - a_1^+ m_1.$$

By the adjunction formula,

$$\deg K_B = \deg(K_{B+}|_B) + \deg N_{B/B+}$$

On the other hand, if we set

$$\begin{aligned} B_i &= B(a_1, \dots, a_i^+, \dots, a_k) \quad i = 1, \dots, \ell \\ &= B(a_1, \dots, \hat{a}_i, \dots) \quad i > \ell \end{aligned}$$

then clearly

$$(11) \quad \deg N_{B/B+} = \sum_{i=1}^k \deg N_{B/B_i},$$

so it suffices to evaluate $\deg N_{B/B_i}$.

Case 1. $i \leq \ell$.

We then have a Cartesian diagram

$$\begin{array}{ccc} s_{A_i} & \longrightarrow & A_i \\ \cap & & \cap \\ s_{A_i^+} & \longrightarrow & A_i^+ \end{array}$$

from which clearly

$$\pi^* N_{B/B_i} = N_{s_{A_i}/s_{A_i^+}} = f^*((a_i - a_i^+) \mathcal{O}(1))$$

hence

$$(12) \quad \deg N_{B/B_i} = (a_i - a_i^+) N(a_1, \dots, a_i + 1, \dots)$$

Case 2. $i > \ell$

We then have a Cartesian diagram

$$\begin{array}{ccccc} & & s_{A_i} & \longrightarrow & A_i \\ & & \cap & & \\ B & \longleftarrow & X & & \cap \\ \downarrow & & \downarrow & & \\ B_i & \longleftarrow & X_i & \longrightarrow & \mathbb{P}^n \end{array}$$

from which as above

$$\deg N_{s_{A_i}/X-i} = a_i N_d(a_1, \dots, a_i + 1, \dots).$$

On the other hand

$$\begin{aligned} \deg N_{s_{A_i}/X_i} &= \deg N_{s_{A_i}/X} + \deg N_{X/X_i}|_{s_{A_i}} \\ &= -m_i + \deg \pi^* N_{B/B_i}. \end{aligned}$$

Consequently

$$(13) \quad \deg N_{B/B_i} = a_i N_d(a_1, \dots, a_i + 1, \dots) + m_i.$$

Putting (9) (10) (11) together, we have computed $\deg(K_B)$, as claimed.

References.

1. Z. Ran : 'Bend, break and count' (alg-geom 9704004)
2. Z. Ran: 'Bend, break and count II: elliptics, cuspidals, linear genera' (alg-geom 9708013)
3. W. Fulton and R. Pandharipande: 'Notes on stable maps and quantum cohomology' Mittag-Leffler Report No. 4 1996/97